



TNFAIP1 contributes to the neurotoxicity induced by Aβ25–35 in Neuro2a cells

Citation

Liu, N., Z. Yu, Y. Xun, M. Li, X. Peng, Y. Xiao, X. Hu, et al. 2016. "TNFAIP1 contributes to the neurotoxicity induced by Aβ25–35 in Neuro2a cells." BMC Neuroscience 17 (1): 51. doi:10.1186/s12868-016-0286-3. <http://dx.doi.org/10.1186/s12868-016-0286-3>.

Published Version

doi:10.1186/s12868-016-0286-3

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:27822171>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

RESEARCH ARTICLE

Open Access



TNFAIP1 contributes to the neurotoxicity induced by A β _{25–35} in Neuro2a cells

Ning Liu^{1,2,3†}, Zhanyang Yu^{3†}, Yu Xun², Miaomiao Li², Xiaoning Peng¹, Ye Xiao², Xiang Hu², Yi Sun⁴, Manjun Yang², Shiquan Gan², Shishan Yuan¹, Xiaoying Wang³, Shuanglin Xiang^{2*} and Jian Zhang^{2*}

Abstract

Background: Amyloid-beta (A β) accumulation is a hallmark of Alzheimer's disease (AD) that can lead to neuronal dysfunction and apoptosis. Tumor necrosis factor, alpha-induced protein 1 (TNFAIP1) is an apoptotic protein that was robustly induced in the transgenic *C. elegans* AD brains. However, the roles of TNFAIP1 in AD have not been investigated.

Results: We found TNFAIP1 protein and mRNA levels were dramatically elevated in primary mouse cortical neurons and Neuro2a (N2a) cells exposed to A β _{25–35}. Knockdown and overexpression of TNFAIP1 significantly attenuated and exacerbated A β _{25–35}-induced neurotoxicity in N2a cells, respectively. Further studies showed that TNFAIP1 knockdown significantly blocked A β _{25–35}-induced cleaved caspase 3, whereas TNFAIP1 overexpression enhanced A β _{25–35}-induced cleaved caspase 3, suggesting that TNFAIP1 plays an important role in A β _{25–35}-induced neuronal apoptosis. Moreover, we observed that TNFAIP1 was capable of inhibiting the levels of phosphorylated Akt and CREB, and also anti-apoptotic protein Bcl-2. TNFAIP1 overexpression enhanced the inhibitory effect of A β _{25–35} on the levels of p-CREB and Bcl-2, while TNFAIP1 knockdown reversed A β _{25–35}-induced attenuation in the levels of p-CREB and Bcl-2.

Conclusion: These results suggested that TNFAIP1 contributes to A β _{25–35}-induced neurotoxicity by attenuating Akt/CREB signaling pathway, and Bcl-2 expression.

Keywords: TNFAIP1, Amyloid-beta, Alzheimer's disease, Neurotoxicity, Neuro2a cells

Background

Alzheimer's disease (AD) is a chronic neurodegenerative disease that is characterized by the accumulation of amyloid-beta (A β) plaques, neurofibrillary tangles, and neuronal loss in various brain regions [1–3]. The 37–43 amino acid A β fragments in the brain are originally derived from the β -amyloid precursor protein (APP) via proteolytic processing by β - and γ -secretase [4]. A β _{1–40} and A β _{1–42} are two major neurotoxic A β fragments, which were mainly generated in the amyloidogenic processing pathway by the action of β - and γ -secretase [4]. A β _{25–35} is not naturally generated in the brain, instead it

represents the neurotoxic fragment of A β _{1–40} or A β _{1–42}, and therefore is often used to mimic the neurotoxic role of A β _{1–40} or A β _{1–42} in experimental studies [5]. Despite increasing evidence showing that A β could cause morphological and biochemical characteristics of apoptosis such as chromatin condensation, DNA fragmentation, caspase activation and subsequent activation of apoptotic signaling pathways [6–8], the molecular mechanisms underlying the neurotoxic effect of A β have not been fully elucidated. Therefore, identification of key players and mechanisms of A β -induced neurotoxicity would benefit the development of novel therapeutic strategies for preventing and treating AD.

Tumor necrosis factor, alpha-induced protein 1 (TNFAIP1) was originally identified as a gene whose expression can be induced by the tumor necrosis factor alpha (TNF α) in umbilical vein endothelial cells [9]. *TNFAIP1* gene is found to be an evolutionarily extremely

*Correspondence: xshlin@hunnu.edu.cn; zhangjian@hunnu.edu.cn

[†]Ning Liu and Zhanyang Yu contributed equally to this work

² Key Laboratory of Protein Chemistry and Development Biology of State Education Ministry of China, College of Life Sciences, Hunan Normal University, Changsha 410081, China

Full list of author information is available at the end of the article

conserved single-copy gene [9], implying that TNFAIP1 has an important physiological role, which is yet to be explored. TNFAIP1 has been demonstrated to interact directly with proliferating cell nuclear antigen (PCNA) and the small subunit (p50) of DNA polymerase δ , implying that it may be involved in DNA synthesis or DNA repair [10, 11]. Kim et al. [12] found that RhoB induces apoptosis by interacting with TNFAIP1 via a JNK-mediated signaling mechanism, suggesting that TNFAIP1 is an apoptosis-related protein. In addition, the transcription levels of TNFAIP1 had been found to be robustly induced in the transgenic *C. elegans* AD brains and post-mortem AD brain [13, 14], suggesting TNFAIP1 may also involve in the process of AD development. Moreover, a recent study implied that estrogen may affect hippocampal-related diseases by regulating TNFAIP1 [15]. However, the role of TNFAIP1 in AD has not been demonstrated.

In the present study, we examined the roles of TNFAIP1 in $A\beta_{25-35}$ -induced apoptosis in neuronal cell line by testing whether the neuronal apoptosis induced by $A\beta_{25-35}$ is associated with the expression of TNFAIP1 protein, and if so, whether apoptosis can be blocked by inhibition of TNFAIP1 expression using TNFAIP1 siRNA. In addition, to further clarify the signal transduction pathways involved in the neurotoxicity induced by $A\beta$, we also examined the potential signal transduction pathways involved in the apoptosis induced by TNFAIP1.

Our findings demonstrated that TNFAIP1 can be induced by $A\beta_{25-35}$. Overexpression of TNFAIP1 promotes $A\beta_{25-35}$ -induced neurotoxicity, whereas knock-down of TNFAIP1 blocks $A\beta_{25-35}$ -induced neurotoxicity. In addition, our results suggested that TNFAIP1 induced by $A\beta_{25-35}$ can further inactivate the Akt/CREB signaling pathway, which in turn downregulates Bcl-2 expression.

Methods

Cell culture and transfection

Animal experiments were following protocols approved by the Ethic Committee of Hunan Normal University, and the Institutional Animal Care and Use Committee of Massachusetts General Hospital in compliance with the NIH Guide for the Care and Use of Laboratory Animals. Primary mouse cortical neurons were isolated from 15 day embryonic cortex obtained from pregnant C57BL/6 female mouse as described before [16]. The cell pellets were resuspended in neuron basal medium, supplemented with 2 % B27 supplement. Cells were seeded at a density of 3×10^5 cells/mL into 6 cm wells plate pre-coated with poly-D-lysine. Medium was half changed every 4 days. All experiments were performed on cultures at days 7–9 in vitro.

The mouse Neuro2A (N2a) neuroblastoma cell line was obtained from the American type culture collection

(ATCC). The N2a cell line was cultured in a humidified (5 % CO_2 , 37 °C) incubator in DMEM supplemented with 10 % heat-inactivated FBS and 50 U/mL penicillin/streptomycin (Invitrogen). The cells were seeded at 1.5×10^5 cells/mL in 24-wells plates or 6-wells plates. For transfection, N2a cells were grown on 24-wells plates to approximately 70 % confluence and then transiently transfected with Control siRNA and TNFAIP1 siRNA (Santa Cruz Biotechnology) or pCMV-myc and pCMV-myc-TNFAIP1 (Myc-TNFAIP1, cloned previously [11]) using Lipofectamine™ 2000 (Invitrogen) following the manufacturer's protocol.

Preparation of β -amyloid fragment $A\beta_{25-35}$

$A\beta_{25-35}$ was purchased from Sigma-Aldrich, and dissolved in deionized distilled water at 1 mM and then aged for 5 days in a humid chamber at 37 °C before being added to the culture medium [17]. For treatment of cells, 1 mM of $A\beta_{25-35}$ was further added into the medium at a final concentration of 2, 5, 10 and 20 μ M, respectively.

Cell viability assay

Cell viability was measured by MTT assay according to manufacturer's protocol. N2a cells were grown on 24-wells plates to approximately 70 % confluence and then transfected with TNFAIP1 siRNA or Myc-TNFAIP1 for 24 h, followed by treatment of $A\beta_{25-35}$ for another 24 h. After treatments, cells were rinsed with PBS and replaced with fresh DMEM containing MTT. The cells were then incubated at 37 °C for 4 h under 5 % CO_2 /95 % air. After the medium was removed, DMSO was added into each well, the absorbance at 570 nm was measured on a microplate reader. Percentages of live cell counts were used for assay normalization.

Real-time PCR

The RT-PCR primers for mouse TNFAIP1 and 18sRNA were purchased from SABiosciences. Total RNA from N2a cells was isolated with RNeasy Lipid Tissue Mini Kit (Qiagen) according to the manufacturer's instructions. Then cDNA was synthesized with SuperScript system (Invitrogen). The mRNA levels of TNFAIP1 and 18sRNA were measured by quantitative RT-PCR using SYBR green kit (Applied Biosystems) in an ABI 7000 real-time PCR system (Applied Biosystems). The PCR conditions were as follows: Initial denaturation of DNA, 95 °C for 5 min; denaturation, 32 cycles of 95 °C for 35 s; annealing, 60 °C for 35 s; extension, 72 °C for 35 s; and final extension, 72 °C for 5 min. Data were analyzed according to the comparative threshold cycle method with expression for sample normalization. RT-PCR assay was performed in triplicate for each sample to ensure reproducibility.

Western blot analysis

Mouse primary cortical neurons or N2a cells were rinsed with PBS, then lysed in cell lysis buffer containing protease inhibitors (Sigma). Equal amounts of total protein (20 µg) were separated using 4–12 % SDS-PAGE gel (Invitrogen) and transferred onto nitrocellulose membrane (Invitrogen). The membrane was blocked with 5 % dry milk in 10 mM PBS buffer (pH 7.2) for 1 h at room temperature. Immunoblots were then performed overnight at 4 °C by incubation with the following primary antibodies: rabbit polyclonal antibodies against CREB (1:2000, Cell Signaling Technology), pCREB (1:1000, Cell Signaling Technology), AKT (1:1000, Cell Signaling Technology), pAKT (1:1000, Cell Signaling Technology), caspase-3 (1:1000, Cell Signaling Technology), cleaved caspase-3 (1:1000, Cell Signaling Technology), TNFAIP1 (1:500, Nanjing Chuanbo Biotech Co, Ltd, China), and mouse monoclonal β -actin antibody (1:5000, sigma), followed by incubation with horseradish peroxidase-conjugated secondary antibodies (1:5000, goat anti-mouse IgG-HRP or goat anti-rabbit IgG-HRP, Abmart) for 1 h at 37 °C. After washed by TTBS, immunolabeling was detected by enhanced chemiluminescence (ECL, Amersham Pharmacia Biotech) according to the manufacturer's protocol and then exposed to film (X-Omat; Eastman Kodak Co.).

Statistical analysis

Data were expressed as mean \pm SD. Three to five separate experiments were performed. Data were analyzed using ANOVA with Tukey post hoc tests (SPSS version 18.0). Statistical significance was at $p < 0.05$.

Results

$A\beta_{25-35}$ induces TNFAIP1 expression

To determine whether $A\beta_{25-35}$ can increase the TNFAIP1 protein expression, mouse primary cortical neurons were treated with $A\beta_{25-35}$ at different doses for 24 h, and Western blot was used to examine TNFAIP1 protein level. As shown in Fig. 1a, TNFAIP1 expression was increased by $A\beta_{25-35}$ in a dose-dependent manner, and 5, 10 and 20 µM of $A\beta_{25-35}$ significantly upregulated TNFAIP1 protein levels (Fig. 1b). Similar results were obtained in N2a cells (Fig. 1c), as TNFAIP1 protein expression in N2a was also significantly increased by $A\beta_{25-35}$ in a dose-dependent manner (Fig. 1d). Furthermore, the endogenous *TNFAIP1* mRNA levels in N2a cells were detected by Real-time PCR. As expected, $A\beta_{25-35}$ treatment resulted in a significant increase in *TNFAIP1* mRNA levels, and treatment with 20 µM of $A\beta_{25-35}$ for 16 h led to the highest expression of *TNFAIP1* mRNA, but prolonged 24 h treatment did not further increase the *TNFAIP1* mRNA levels (Fig. 1e).

TNFAIP1 contributes to $A\beta_{25-35}$ -induced cell toxicity in N2a cells

To further investigate whether TNFAIP1 is involved in $A\beta_{25-35}$ -induced cell death, N2a cells were transiently transfected with Control siRNA or TNFAIP1 siRNA for 24 h, and then treated with indicated dose of $A\beta_{25-35}$ for another 24 h. Western blot showed that the endogenous TNFAIP1 protein expression was significantly suppressed by specific TNFAIP1 siRNA but not Control siRNA (Fig. 2a). MTT assay indicated that the cell viability in Control siRNA transfected-N2a cells was significantly decreased when exposed to 10 or 20 µM of $A\beta_{25-35}$. However, inhibition of TNFAIP1 by TNFAIP1 siRNA significantly rescued the viability of N2a cells under 20 µM $A\beta_{25-35}$ (Fig. 2b). Furthermore, to investigate whether TNFAIP1 is associated with increased susceptibility to $A\beta_{25-35}$, the effect of TNFAIP1 overexpression on $A\beta_{25-35}$ -reduced the cell viability was also investigated. Transfection of Myc-TNFAIP1 into N2a cells led to significant overexpression of TNFAIP1 protein (Fig. 2c), and $A\beta_{25-35}$ -induced cell death was further significantly exacerbated by TNFAIP1 overexpression (Fig. 2d). Taken together, these results indicate that TNFAIP1 contributes to $A\beta_{25-35}$ -induced cell toxicity of N2a cells.

TNFAIP1 contributes to $A\beta_{25-35}$ -induced apoptosis

Previous study suggested that TNFAIP1 is an apoptosis-related protein [12], implying that TNFAIP1 may also involve in $A\beta_{25-35}$ -induced apoptosis. Cleaved caspase-3 is a pivotal executioner and hallmark of apoptosis that is usually activated in the apoptotic cell by both extrinsic and intrinsic pathways. To further determine whether involvement of TNFAIP1 in $A\beta_{25-35}$ -induced neuronal apoptosis is associated with changes in the activities of caspases, cleaved caspase-3 was detected by Western blot. As shown in Fig. 3a, b, N2a cells treated with 20 µM of $A\beta_{25-35}$ led to a significant increase in the level of cleaved caspase-3, while $A\beta_{25-35}$ -induced caspase-3 cleavage was significantly repressed by the knockdown of TNFAIP1. In contrast, 20 µM of $A\beta_{25-35}$ treatment or Myc-TNFAIP1 transfection or $A\beta_{25-35}$ +Myc-TNFAIP1 co-treatment exhibited a significant increase in the cleavage of caspase-3 (Fig. 3c, d). These results indicate that TNFAIP1 contributes to $A\beta_{25-35}$ -induced apoptosis.

TNFAIP1 inhibits Akt/CREB signaling pathway

It has been reported that Akt/CREB signaling plays neuroprotective roles [18] and $A\beta$ could downregulate Akt survival pathway [19]. In agreement with previous studies, our results showed that $A\beta_{25-35}$ causes a decrease in Akt phosphorylation (phosphor-Ser473), CREB phosphorylation (phosphor-Ser133-CREB) and




Fig. 1 Effect of Aβ₂₅₋₃₅ on TNFAIP1 expression. **a** TNFAIP1 protein levels were measured by Western blot in mouse primary cortical neurons cultured in 6 wells plate and treated with dose-dependent Aβ₂₅₋₃₅ (0–20 μM). **b** Quantitative analysis of TNFAIP1 protein levels (fold of control) in mouse primary cortical neurons. Data are expressed as mean ± SD, n = 4, *p < 0.05 compared with Aβ₂₅₋₃₅ (0 μM) (PBS). **c** TNFAIP1 protein levels were measured by Western blot in N2a cells cultured in 6 wells plate and treated with dose-dependent Aβ₂₅₋₃₅ (0–20 μM). **d** Quantitative analysis of TNFAIP1 protein levels (fold of control) in N2a cells. Data are expressed as mean ± SD, n = 4, *p < 0.05 compared with Aβ₂₅₋₃₅ (0 μM) (PBS). **e** TNFAIP1 mRNA levels were determined by RT-PCR in N2a cells cultured in 6 wells plate and treated with 20 μM of Aβ₂₅₋₃₅. Data are expressed as mean ± SD, n = 3, *p < 0.05 compared with Aβ₂₅₋₃₅ (0 h)

Bcl-2 expression in a dose-dependent manner (Fig. 4a). To further explore whether involvement of TNFAIP1 in neuronal apoptosis was associated with changes in Akt/CREB signaling pathway, we overexpressed TNFAIP1 in N2a cells in dose-dependent manner and then assessed Akt phosphorylation and CREB phosphorylation (Fig. 4a). As expected, overexpression of TNFAIP1 in N2a cells led to significant attenuation of p-Akt and p-CREB in a dose-dependent manner (Fig. 4b–d). Importantly, we also observed that overexpression of TNFAIP1 could significantly reduce the protein levels of Bcl-2 (Fig. 4e), which is an important neuroprotectant and transactivated by CREB in response to Aβ stimulation [20]. Therefore, these results indicate that TNFAIP1 may

mediate Aβ₂₅₋₃₅-induced apoptotic signaling by attenuating p-Akt, p-CREB and Bcl-2 expression.

TNFAIP1 knockdown recovers Aβ₂₅₋₃₅-induced CREB dephosphorylation and Bcl-2 downregulation

Aβ had been reported to dramatically decrease CREB phosphorylation and Bcl-2 expression [20, 21]. To investigate the roles of TNFAIP1 in Aβ₂₅₋₃₅-reduced CREB phosphorylation and Bcl-2 expression, control siRNA or TNFAIP1 siRNA was transfected into N2a cells for 24 h, followed by treatment with 20 μM of Aβ₂₅₋₃₅. Consistent with previous studies [20], Aβ₂₅₋₃₅ treatment significantly reduced CREB phosphorylation and Bcl-2 expression. TNFAIP1 siRNA but not control siRNA significantly




Fig. 2 Effect of TNFAIP1 on Aβ₂₅₋₃₅-induced neuronal cell death. **a** TNFAIP1 protein levels were measured by Western blot in N2a cells cultured in 12 wells plate and transfected with 20 pmol of Control siRNA or TNFAIP1 siRNA. **b** MTT reduction was measured by MTT assay in N2a cells cultured in 12 wells plate and transfected with 20 pmol of Control siRNA or TNFAIP1 siRNA, followed by treated by dose-dependent Aβ₂₅₋₃₅ (0–20 μM). Data are expressed as mean ± SD, n = 4, *p < 0.05 compared with Aβ₂₅₋₃₅ (0 μM), #p < 0.05 compared with Control siRNA plus Aβ₂₅₋₃₅ (20 μM). **c** Myc-TNFAIP1 protein levels were measured by Western blot in N2a cells cultured in 12 wells plate and transfected with 1.2 μg of pCMV-Myc or Myc-TNFAIP1. **d** MTT reduction was measured by MTT assay in N2a cells cultured in 12 wells plate and transfected with dose-dependent Myc-TNFAIP1 (0–1.2 μg), followed by treated by PBS or Aβ₂₅₋₃₅ (20 μM). Data are expressed as mean ± SD, n = 4, *p < 0.05 compared with Myc-TNFAIP1 (0 μg) plus PBS. #p < 0.05 compared with Myc-TNFAIP1 plus PBS

upregulated CREB phosphorylation. However, TNFAIP1 siRNA could partially attenuate the inhibitory effect of Aβ₂₅₋₃₅ on p-CREB and Bcl-2 (Fig. 5). These results indicated that inhibition of TNFAIP1 could attenuate Aβ-induced neurotoxicity partially by recovering the inhibitory effect of Aβ on p-CREB and Bcl-2 expression.

TNFAIP1 contributes to Aβ₂₅₋₃₅-induced CREB dephosphorylation and Bcl-2 downregulation

The effect of TNFAIP1 overexpression on CREB phosphorylation and Bcl-2 expression under resting and Aβ₂₅₋₃₅ treatment condition were also determined. Our results showed that 20 μM of Aβ₂₅₋₃₅ or Aβ₂₅₋₃₅+Myc-TNFAIP1 had exhibited almost equivalent inhibitory effect on the p-CREB (Fig. 6a, b). Moreover, we also observed that 20 μM of Aβ₂₅₋₃₅ treatment, Myc-TNFAIP1 transfection, and Aβ₂₅₋₃₅+Myc-TNFAIP1 co-treatment result in significant decrease in the expression of Bcl-2 (Fig. 6a, c). These results implied that TNFAIP1 maybe a key

mediator involved in Aβ-induced neurotoxicity by inhibition of p-CREB and Bcl-2 expression.

Discussion

Deposition of Aβ in the brain is a pathological hallmark of AD as it is responsible for progressive neurodegeneration in AD [22]. Accumulating evidences indicated that Aβ could cause neurotoxicity by inducing neuronal apoptosis in vitro and in vivo [23, 24]. Elucidating the molecular mechanisms underlying Aβ-induced neuronal apoptosis will help us on developing treatment of AD. However, despite intense research, it remains unclear how Aβ triggers the signaling cascade that results in neuronal dysfunction and neurotoxicity. In the present study, our results demonstrated that TNFAIP1 protein is induced by Aβ₂₅₋₃₅ and involved in Aβ₂₅₋₃₅-induced neuronal apoptotic pathway because overexpression of TNFAIP1 can accelerate Aβ₂₅₋₃₅-induced apoptosis while inhibition of TNFAIP1 can significantly reduce




Fig. 3 Effect of TNFAIP1 on Aβ₂₅₋₃₅-induced the cleavage of caspase-3. **a** Cleaved caspase-3 protein levels were measured by Western blot in N2a cells cultured in 6 wells plate and transfected with 50 pmol of Control siRNA or TNFAIP1 siRNA, followed by treated by Aβ₂₅₋₃₅ (20 μM). **b** Quantitative analysis of Cleaved caspase-3 protein levels (fold of control) in N2a cells. Data are expressed as mean ± SD, n = 3, *p < 0.05 compared with Control siRNA, #p < 0.05 compared with Control siRNA plus Aβ₂₅₋₃₅ (20 μM). **c** Cleaved caspase-3 protein levels were measured by Western blot in N2a cells cultured in 6 wells plate and transfected with 4 μg of pCMV-Myc or Myc-TNFAIP1, followed by treated by Aβ₂₅₋₃₅ (20 μM). **d** Quantitative analysis of Cleaved caspase-3 protein levels (fold of control) in N2a cells. Data are expressed as mean ± SD, n = 3, *p < 0.05, **p < 0.01 compared with pCMV-Myc

Aβ₂₅₋₃₅-induced neuronal apoptosis. Moreover, we found that TNFAIP1 was involved in Aβ₂₅₋₃₅-induced neuronal apoptosis by deactivating Akt signaling pathway and CREB transcriptional activity.

The transcript levels of TNFAIP1 was found to be robustly induced in the transgenic *C. elegans* AD brains and post-mortem AD brain in previous study [12, 13], which led us to examine the hypothesis that TNFAIP1 was involved in the pathological development of AD using an in vitro mouse AD model: mouse primary cortical neurons and N2a neuroblastoma cells treated by Aβ₂₅₋₃₅. Our results suggested that Aβ₂₅₋₃₅ could induce TNFAIP1 protein and mRNA levels in a dose-dependent manner. Furthermore, overexpression or knockdown of TNFAIP1 exacerbated or alleviated Aβ₂₅₋₃₅-induced neurotoxicity. Together, these results indicated that the TNFAIP1 gene expression may be critical for Aβ₂₅₋₃₅-induced neuronal death and intervention of its expression could be a potential method for inhibiting Aβ₂₅₋₃₅-induced neurotoxicity. However, the regulatory mechanisms of TNFAIP1 remain largely unclear. Only a recent report described that transcription factor Sp1 could bind to human *TNFAIP1* promoter region and

transactivate *TNFAIP1* promoter [25]. As TNFAIP1 had been originally found to be induced by TNFα [9], which is a potent activator of NFκB, thus it is possible that TNFAIP1 was also regulated by NFκB when exposed to Aβ₂₅₋₃₅. Further studies will be performed to elucidate whether inhibition of NFκB activation can reduce Aβ₂₅₋₃₅-induced *TNFAIP1* gene expression.

Previous studies described that TNFAIP1 was a pro-apoptotic protein [12], and apoptosis is a general neuronal death pathway in neurodegenerative diseases which could be triggered by exposure to Aβ. In the present study, we extended our observations to in vitro AD models that TNFAIP1 was also involved in Aβ₂₅₋₃₅-induced apoptosis. We found that inhibition of *TNFAIP1* gene by specific siRNA significantly attenuated Aβ₂₅₋₃₅ induced cleaved-caspase 3. As caspase-3 cleavage is a central event in executing Aβ₂₅₋₃₅-induced neuronal apoptosis [26, 27], these results provide further evidence that TNFAIP1 was involved in Aβ₂₅₋₃₅-induced apoptosis. However, it is worth noting that Aβ₂₅₋₃₅+Myc-TNFAIP1 co-treatment did not cause a much higher increase in the cleavage of caspase-3 than Aβ₂₅₋₃₅ treatment or Myc-TNFAIP1 transfection alone. It is possible that 20 μM of






Fig. 4 Effect of Aβ₂₅₋₃₅ and TNFAIP1 on p-Akt, p-CREB and Bcl-2 expression. **a** p-Akt, p-CREB and Bcl-2 protein levels were measured by Western blot in N2a cells cultured in 6 wells plate and treated with dose-dependent Aβ₂₅₋₃₅ (0–20 μM). **b** p-Akt, p-CREB and Bcl-2 protein levels were measured by Western blot in N2a cells cultured in 6 wells plate and transfected with Myc-TNFAIP1 and pCMV-Myc (0–4 μg). **c** Quantitative analysis of p-CREB protein levels (fold of control). Data are expressed as mean ± SD, n = 3, *p < 0.05, **p < 0.01 compared with Myc-TNFAIP1 (0 μg). **d** Quantitative analysis of p-Akt protein levels (fold of control). Data are expressed as mean ± SD, n = 3, *p < 0.05 compared with Myc-TNFAIP1 (0 μg). **e** Quantitative analysis of Bcl-2 protein levels (fold of control). Data are expressed as mean ± SD, n = 3, *p < 0.05 compared with Myc-TNFAIP1 (0 μg)

Aβ₂₅₋₃₅ treatment or Myc-TNFAIP1 transfection alone had already exerted a relatively strong apoptotic effect on N2a cells, which led to a insignificant effect on caspase 3 cleavage by additional other stimuli. This issue will be further defined in future studies. Thus, these results are in support of the hypothesis that the induction of TNFAIP1 expression is part of the intrinsic program of apoptotic neuronal death induced by accumulation of Aβ in AD.

It had been demonstrated that CREB is directly phosphorylated and regulated by the protein kinase Akt [28], and Akt/CREB signal has been shown to play a pivotal role in neuroprotection by enhancing cell survival and inhibiting apoptosis [18]. Importantly, Akt and CREB could promote neuronal cell survival by upregulating Bcl-2 protein levels [29]. As expected, TNFAIP1 could

reduce p-Akt, p-CREB and Bcl-2 protein levels in a dose-dependent manner. Moreover, we observed overexpression of TNFAIP1 could further attenuate Aβ-inhibited p-CREB and Bcl-2 protein expression, whereas inhibition of TNFAIP1 could significantly reverse the effect of Aβ on p-CREB and Bcl-2. These results suggest that TNFAIP1 may mediate the deactivation of CREB induced by Aβ, and then trigger neuronal apoptosis. MAPKs had been reported to be involved in Aβ-induced apoptosis, and MAPKs signaling cascades were activated in brains from AD patients [30]. Moreover, Aβ-induced MAPKs include ERK1/2, JNK and p38 MAPK had been demonstrated to be associated with neuronal cell death [31, 32]. Therefore, TNFAIP1 may also modulate MAPKs signaling cascades involved in Aβ-induced neuronal apoptosis. Further study will be conducted to elucidate the role



of TNFAIP1 in modulating MAPKs signaling pathway involved in the apoptotic effect of A β .

In summary, in this study we found that TNFAIP1 expression was elevated in mouse primary cortical neurons and N2a cells exposed to A β_{25-35} . TNFAIP1 overexpression was correlated with neuronal apoptosis

initiated by A β_{25-35} and inhibition of TNFAIP1 was able to reduce the neuronal damage induced by A β_{25-35} in vitro AD models. Moreover, our results suggested that TNFAIP1 may mediate A β_{25-35} -induced neuronal damage by inhibiting p-CREB and Bcl-2 protein levels. Our results implied that inhibition of *TNFAIP1* gene during

AD process could be a potential therapeutic strategy for treatment of AD. Further studies will be performed to determine the roles of TNFAIP1 in in vivo AD models.

Conclusions

In summary, this study clearly demonstrated that TNFAIP1 was significantly upregulated by A β _{25–35} in mouse primary cortical neurons and N2a cells, and TNFAIP1 may be a key player that mediated A β _{25–35}-induced neurotoxicity by inactivating the Akt/CREB signaling pathway, and in turn downregulating anti-apoptotic protein Bcl-2. Our findings implied that TNFAIP1 may be a potential therapeutic target for treatment of AD, but the roles and mechanism of TNFAIP1 in in vivo AD models need to be further elucidated in our future study.

Abbreviations

A β : amyloid-beta; AD: Alzheimer's disease; N2a: Neuro2a; TNFAIP1: tumor necrosis factor, alpha-induced protein 1; TNF α : tumor necrosis factor alpha; APP: β -amyloid precursor protein; PCNA: proliferating cell nuclear antigen; ATCC: The American type culture collection; RT-PCR: real-time PCR.

Authors' contributions

Designed the experiments NL, ZY, XW, SX, and JZ. Performed the experiments NL, ZY, YX, ML, YX, XH, YS, MY, SG, SY and XP. Analyzed the data NL, ZY, and YX. Wrote the paper NL, ZY, XW, SX, and JZ. All authors read and approved the final manuscript.

Author details

¹ College of Medicine, Hunan Normal University, Changsha, China. ² Key Laboratory of Protein Chemistry and Development Biology of State Education Ministry of China, College of Life Sciences, Hunan Normal University, Changsha 410081, China. ³ Neuroprotection Research Laboratory, Department of Neurology and Radiology, Massachusetts General Hospital, Neuroscience Program, Harvard Medical School, Boston, MA, USA. ⁴ Department of Pathology, The Second Xiangya Hospital of Central South University, Changsha, Hunan, China.

Acknowledgements

This work was partially supported by the financial support from the program of China Scholarships Council (No. 2009672005), the China Postdoctoral Science Foundation (2014M552139), Scientific Research Fund of Hunan Provincial Education Department (2014RS4008), and AHA FDA Summer 2011 Postdoctoral Fellowship (12POST9720007).

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

All the data supporting our findings is contained within the manuscript.

Ethics

Animal experiments were following protocols approved by the Ethic Committee of Hunan Normal University, and the Institutional Animal Care and Use Committee of Massachusetts General Hospital in compliance with the NIH Guide for the Care and Use of Laboratory Animals.

Funding

This work was supported in part by the financial support from the program of China Scholarships Council (No. 2009672005), the China Postdoctoral Science Foundation (2014M552139), Scientific Research Fund of Hunan Provincial Education Department (2014RS4008) (to N. Liu), and AHA FDA Summer 2011 Postdoctoral Fellowship (12POST9720007) (to Z. Yu). The funders had no role

in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Received: 7 October 2015 Accepted: 8 July 2016

Published online: 18 July 2016

Reference

- Selkoe DJ. The molecular pathology of Alzheimer's disease. *Neuron*. 1991;6(4):487–98.
- Hardy JA, Higgins GA. Alzheimer's disease: the amyloid cascade hypothesis. *Science*. 1992;256(5054):184–5.
- Castillo-Carranza DL, Guerrero-Munoz MJ, Sengupta U, Hernandez C, Barrett AD, Dineley K, Kaye R. Tau immunotherapy modulates both pathological tau and upstream amyloid pathology in an Alzheimer's disease mouse model. *J Neurosci*. 2015;35(12):4857–68.
- Haass C, Kaether C, Thinakaran G, Sisodia S. Trafficking and proteolytic processing of APP. *Cold Spring Harb Perspect Med*. 2012;2(5):a006270.
- Kaminsky YG, Marlatt MW, Smith MA, Kosenko EA. Subcellular and metabolic examination of amyloid-beta peptides in Alzheimer disease pathogenesis: evidence for A β 25–35. *Exp Neurol*. 2010;221(1):26–37.
- Mattson MP. Apoptosis in neurodegenerative disorders. *Nat Rev Mol Cell Biol*. 2000;1(2):120–9.
- Friedlander RM. Apoptosis and caspases in neurodegenerative diseases. *N Engl J Med*. 2003;348(14):1365–75.
- Dineley KT, Westerman M, Bui D, Bell K, Ashe KH, Sweatt JD. Beta-amyloid activates the mitogen-activated protein kinase cascade via hippocampal α 7 nicotinic acetylcholine receptors: In vitro and in vivo mechanisms related to Alzheimer's disease. *J Neurosci*. 2001;21(12):4125–33.
- Wolf FW, Marks RM, Sarma V, Byers MG, Katz RW, Shows TB, Dixit VM. Characterization of a novel tumor necrosis factor- α -induced endothelial primary response gene. *J Biol Chem*. 1992;267(2):1317–26.
- Zhou J, Hu X, Xiong X, Liu X, Liu Y, Ren K, Jiang T, Zhang J. Cloning of two rat PDIP1 related genes and their interactions with proliferating cell nuclear antigen. *J Exp Zool A Comp Exp Biol*. 2005;303(3):227–40.
- Yang L, Liu N, Hu X, Zhang W, Wang T, Li H, Zhang B, Xiang S, Zhou J, Zhang J. CK2 phosphorylates TNFAIP1 to affect its subcellular localization and interaction with PCNA. *Mol Biol Rep*. 2010;37(6):2967–73.
- Kim DM, Chung KS, Choi SJ, Jung YJ, Park SK, Han GH, Ha JS, Song KB, Choi NS, Kim HM, et al. RoB induces apoptosis via direct interaction with TNFAIP1 in HeLa cells. *Int J Cancer*. 2009;125(11):2520–7.
- Link CD, Taft A, Kapulkin V, Duke K, Kim S, Fei Q, Wood DE, Sahagan BG. Gene expression analysis in a transgenic *Caenorhabditis elegans* Alzheimer's disease model. *Neurobiol Aging*. 2003;24(3):397–413.
- Wu Y, Luo Y. Transgenic *C. elegans* as a model in Alzheimer's research. *Curr Alzheimer Res*. 2005;2(1):37–45.
- Liu H, Yang L, Zhao Y, Zeng G, Wu Y, Chen Y, Zhang J, Zeng Q. Estrogen is a novel regulator of Tnfaip1 in mouse hippocampus. *Int J Mol Med*. 2014;34(1):219–27.
- Yu Z, Liu N, Li Y, Xu J, Wang X. Neuroglobin overexpression inhibits oxygen-glucose deprivation-induced mitochondrial permeability transition pore opening in primary cultured mouse cortical neurons. *Neurobiol Dis*. 2013;56:95–103.
- Yu MS, Suen KC, Kwok NS, So KF, Hugon J, Chang RC. Beta-amyloid peptides induces neuronal apoptosis via a mechanism independent of unfolded protein responses. *Apoptosis*. 2006;11(5):687–700.
- Carlioni S, Girelli S, Scopa C, Buonocore G, Longini M, Balduini W. Activation of autophagy and Akt/CREB signaling play an equivalent role in the neuroprotective effect of rapamycin in neonatal hypoxia-ischemia. *Autophagy*. 2010;6(3):366–77.
- Magrane J, Rosen KM, Smith RC, Walsh K, Gouras GK, Querfurth HW. Intraneuronal beta-amyloid expression downregulates the Akt survival pathway and blunts the stress response. *J Neurosci*. 2005;25(47):10960–9.
- Wei G, Chen YB, Chen DF, Lai XP, Liu DH, Deng RD, Zhou JH, Zhang SX, Li YW, Lii H, et al. Beta-amyloid inhibits neuronal apoptosis via the CaMKII/CREB/Bcl-2 signaling pathway in an in vitro model and A β PP/PS1 mice. *J Alzheimers Dis*. 2013;33(3):863–80.

21. Tong L, Thornton PL, Balazs R, Cotman CW. Beta-amyloid-(1–42) impairs activity-dependent cAMP-response element-binding protein signaling in neurons at concentrations in which cell survival is not compromised. *J Biol Chem*. 2001;276(20):17301–6.
22. Selkoe DJ. Alzheimer's disease is a synaptic failure. *Science*. 2002;298(5594):789–91.
23. Cancino GI, Toledo EM, Leal NR, Hernandez DE, Yevenes LF, Inestrosa NC, Alvarez AR. ST1571 prevents apoptosis, tau phosphorylation and behavioural impairments induced by Alzheimer's beta-amyloid deposits. *Brain*. 2008;131(Pt 9):2425–42.
24. Loo DT, Copani A, Pike CJ, Whittemore ER, Walencewicz AJ, Cotman CW. Apoptosis is induced by beta-amyloid in cultured central nervous system neurons. *Proc Natl Acad Sci USA*. 1993;90(17):7951–5.
25. Liu M, Sun Z, Zhou A, Li H, Yang L, Zhou C, Liu R, Hu X, Zhou J, Xiang S, et al. Functional characterization of the promoter region of human TNFAIP1 gene. *Mol Biol Rep*. 2010;37(4):1699–705.
26. Harada J, Sugimoto M. Activation of caspase-3 in beta-amyloid-induced apoptosis of cultured rat cortical neurons. *Brain Res*. 1999;842(2):311–23.
27. Marin N, Romero B, Bosch-Morell F, Llansola M, Felipe V, Roma J, Romero FJ. Beta-amyloid-induced activation of caspase-3 in primary cultures of rat neurons. *Mech Ageing Dev*. 2000;119(1–2):63–7.
28. Du K, Montminy M. CREB is a regulatory target for the protein kinase Akt/PKB. *J Biol Chem*. 1998;273(49):32377–9.
29. Ji L, Mochon E, Arcinas M, Boxer LM. CREB proteins function as positive regulators of the translocated bcl-2 allele in t(14;18) lymphomas. *J Biol Chem*. 1996;271(37):22687–91.
30. Swatton JE, Sellers LA, Faull RL, Holland A, Iritani S, Bahn S. Increased MAP kinase activity in Alzheimer's and Down syndrome but not in schizophrenia human brain. *Eur J Neurosci*. 2004;19(10):2711–9.
31. Longpre F, Garneau P, Christen Y, Ramassamy C. Protection by EGb 761 against beta-amyloid-induced neurotoxicity: involvement of NF-kappaB, SIRT1, and MAPKs pathways and inhibition of amyloid fibril formation. *Free Radic Biol Med*. 2006;41(12):1781–94.
32. Behrens MM, Strasser U, Koh JY, Gwag BJ, Choi DW. Prevention of neuronal apoptosis by phorbol ester-induced activation of protein kinase C: blockade of p38 mitogen-activated protein kinase. *Neuroscience*. 1999;94(3):917–27.

Submit your next manuscript to BioMed Central
and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

